


Bioactivity – Antibody Binding


Mouse TrkC-His-Avi dimer, ELISA

0.2 µg of TrkC protein dimer per well



Immobilized mouse TrkC protein dimer, His-Avi Tag (CSP-25176-03) at 2 µg/mL (100 µL/well) can bind anti-mouse TrkC monoclonal antibody with half maximal effective concentration (EC₅₀) range of 1.6-6.4 µg/mL (QC tested).

SDS-PAGE



MW: Molecular Weight marker reduced condition
NR: TrkC dimer under non-reduced condition

The migration range of the heterodimer protein with glycosylation under non-reducing condition is >190 kDa on SDS PAGE.



Mouse TrkC Protein Dimer, His-Avi Tag
Product Code: CSP-25176-03
For Research Use Only (RUO)

Expression Host
HEK293T

Purity
Greater than 90% dimer form as determined by SDS-PAGE under non-reducing condition

Protein Construct
TrkC protein dimer contains the TrkC extracellular domain (UniProt# Q6VNS1) fused with a proprietary cis-dimer motif followed by a tandem His-Avi tag at the C-terminus. Expressed in HEK293T cell line.

SDS-Page Molecular Weight
109 kDa. The migration range of the heterodimer protein with glycosylation under non-reducing condition is >190 kDa on SDS PAGE.

Shipping Conditions
Frozen Dry Ice

Protein Name
TrkC (NTRK3)

Alternate Name(s)
neurotrophic receptor tyrosine kinase 3, NTRK3, NT-3 growth factor receptor, neurotrophic tyrosine kinase receptor type 3, TrkC tyrosine kinase, gp145, GP145-TrkC, TRKC

Amino Acid Range
1: C32-T429

Formulation
0.22µm filtered PBS, pH 7.4

Stability & Storage
-80°C

Background

Tropomyosin receptor kinase C (TrkC) is part of the family of receptor tyrosine kinases and primarily the receptor for neurotrophin-3 (NT-3). TrkC is also known as neurotrophic receptor tyrosine kinase 3 (NTRK3), NT-3 growth factor receptor, neurotrophic tyrosine kinase receptor type 3, gp145, and GP145-TrkC. TrkC is a Type I transmembrane protein. It contains an extracellular domain with two cysteine-rich clusters (C1 and C2), three leucine-rich 24-residue repeats (LRR1–3), and two immunoglobulin-like domains (Ig1 and Ig2) followed by a transmembrane domain and a cytoplasmic domain. TrkC is a high affinity receptor for NT-3, and ligand binding induces receptor dimerization which is required for trans-autophosphorylation that triggers downstream signaling cascades. TrkC plays a role in many cancers, therefore, a recombinant protein mimicking the dimer conformation can be crucial for cancer therapeutic discovery. While structurally and functionally similar to human TrkC homodimer, mouse TrkC homodimer is a species-specific tool essential for preclinical studies, basic research, and translational research.