



## Bioactivity – Antibody Binding

### Human CXCR4 full length, ELISA 0.5 µg of CXCR4 per well



Immobilized human CXCR4 VLP (Cat. No. CMP-24005) at 5 µg/mL (100 µL/well) can bind anti-human CXCR4 monoclonal antibody with half maximal effective concentration (EC<sub>50</sub>) range of 2.3-9.19 ng/mL (QC tested).

## DLS



The mean peak diameter (nm) of the CMP is 185.3 with a range of 161.3-203.5 with more than 95% intensity as determined by dynamic light scattering (DLS).



## Human CXCR4 Full Length Protein, Displayed on Nanoparticles

Product Code: CMP-24005  
For Research Use Only (RUO)

**Expression Host**

HEK293T

**Protein Name**

Human CXCR4

**Alternate Name(s)**

FB22, HM89, LAP3, LCR1, NPYR, WHIM, CD184, LAP-3, LESTR, NPY3R, NPYRL, WHIMS, HSY3RR, NPYY3R, WHIMS1, and D2S201E

**Amino Acid Range**

Met1-Ser352

**Protein Construct**

The human CXCR4 full length protein (UniProt# P61073), displayed on nanoparticles and mimics the native protein conformation on the cell surface. Expressed in HEK293T cell line.

**Formulation**

0.2µm filtered PBS, pH 7.4

**Shipping Conditions**

Frozen Dry Ice

**Stability & Storage**

-80°C

**Background**

C-X-C chemokine receptor type 4 (CXCR4) is also known as CD184, D2S201E, FB22, HM89, HSY3RR, LAP3, LCR1, LESTR, NPY3R, NPYR, NPYRL, NPYY3R or WHIM. CXCR-4 is a G protein-coupled receptor (GPCR) for the C-X-C motif chemokine 12 (CXCL12), also known as stromal cell-derived factor 1 (SDF-1). CXCR4 also acts as a co-receptor for HIV infection.